

Perturbative matching of heavy-light currents at one-loop

Aida X. El-Khadra* and Elvira Gámiz

Department of Physics, University of Illinois, Urbana, IL 61801, USA

E-mail: axk@uiuc.edu, megamiz@uiuc.edu

Andreas S. Kronfeld

Theoretical Physics Department, Fermi National Accelerator Laboratory, Batavia, IL 60510, USA

E-mail: ask@fnal.gov

Matthew A. Nobes

Department of Physics, Cornell University, Ithaca, NY 14853, USA

and

Brevan Howard Asset Management, London, SW1Y 6XA, United Kingdom

E-mail: drnobes@gmail.com

We present results of a perturbative matching calculation performed at one-loop for heavy-light currents. We use the Fermilab action for the heavy quarks, the Asqtad action for the light quarks, and an improved gluon action. We also present results for heavy-heavy currents with Fermilab heavy quarks and improved glue.

The XXV International Symposium on Lattice Field Theory

July 30-4 August 2007

Regensburg, Germany

*Speaker.

1. Introduction

The Fermilab Lattice and MILC collaboration's program includes calculations of the hadronic matrix elements for weak D and B meson decays, in particular, the decay constants f_D , f_{D_s} , f_B , and f_{B_s} and the semileptonic form factors for $B \rightarrow \pi \ell \nu$, $D \rightarrow \pi(K) \ell \nu$, and $B \rightarrow D^* \ell \nu$. In this work we present a perturbative matching calculation of the relevant current renormalizations to one-loop order. The numerical simulations for the above physics analyses use MILC ensembles with improved glue and $2 + 1$ Asqtad staggered sea quarks [1]. The light valence quarks are also generated from Asqtad staggered quarks and converted to naive quarks. The heavy (charm and beauty) quarks are treated with the Fermilab action. See Ref. [2] for more details on the actions and parameters used in the numerical simulations.

2. Definitions

In this work we follow the analysis of Ref. [3], where the one-loop corrections to heavy-light and heavy-heavy current renormalizations were calculated for Fermilab heavy and Clover light quarks with Wilson glue.

The heavy-light currents have the form

$$J_\mu^{hl\text{lat}} = \bar{\Psi}_h \Gamma_\mu \psi_l, \quad (2.1)$$

where $\Gamma_\mu = \gamma_\mu$ or $\gamma_\mu \gamma_5$ and ψ_l denotes a naive Asqtad Dirac spinor. The Fermilab Dirac spinor, ψ_h , is rotated by

$$\psi_h = \psi [1 + ad_1 \gamma \cdot \mathbf{D}], \quad (2.2)$$

with the tree-level coefficient

$$d_1 = \frac{1}{2 + m_0 a} - \frac{1}{2(1 + m_0 a)}. \quad (2.3)$$

The heavy-heavy currents have the form

$$J_\mu^{hh'\text{lat}} = \bar{\Psi}_h \Gamma_\mu \psi_{h'}, \quad (2.4)$$

where now both spinors are rotated Dirac spinors. Since the heavy quarks are rotated, the lattice currents of Eqns. (2.1) and (2.4) include the leading order tree-level discretization corrections.

The current renormalization is defined as

$$Z_{J_\Gamma}^{hl} = \frac{(Z_{2h}^{(1/2)} \Lambda_{J_\Gamma} Z_{2l}^{(1/2)})^{\text{cont}}}{(Z_{2h}^{(1/2)} \Lambda_{J_\Gamma} Z_{2l}^{(1/2)})^{\text{lat}}}, \quad (2.5)$$

where Λ_{J_Γ} are the vertex corrections and Z_{2h} (Z_{2l}) are the heavy (light) quark wave function renormalizations.

We factor out the dominant mass dependence due to the tree-level wave function renormalization of the heavy Fermilab quark by defining the perturbative expansion as

$$e^{-m_1^{[0]} a/2} Z_{J_\Gamma}^{hl} = 1 + g_0^2 Z_{J_\Gamma}^{hl[1]} + \dots, \quad (2.6)$$

where the heavy quark masses are defined as usual,

$$m_1^{[0]} a = \log(1 + m_0 a), \quad m_0 a = 1/(2\kappa_h) - 1/(2\kappa_{crit}). \quad (2.7)$$

Since Z_{V_4} for degenerate masses is easy to calculate nonperturbatively, it is useful to define

$$\rho_{J_\Gamma}^{hl} \equiv \frac{Z_{J_\Gamma}^{hl}}{\sqrt{Z_{V_4}^{hh} Z_{V_4}^{ll}}} = 1 + g_0^2 \rho_{J_\Gamma}^{[1]} + \dots \quad (2.8)$$

In this case, the dominant mass dependence cancels by construction.

Analogously, for heavy-heavy currents we have:

$$Z_{J_\Gamma}^{hh'} = \frac{(Z_{2h}^{(1/2)} \Lambda_{J_\Gamma} Z_{2h'}^{(1/2)})^{\text{cont}}}{(Z_{2h}^{(1/2)} \Lambda_{J_\Gamma} Z_{2h'}^{(1/2)})^{\text{lat}}}. \quad (2.9)$$

Taking the leading mass dependence out again, the perturbative expansion is defined as

$$e^{-(m_{1h}^{[0]} + m_{1h'}^{[0]})a/2} Z_{J_\Gamma}^{hh'} = 1 + g_0^2 \rho_{J_\Gamma}^{hh'[1]} + \dots \quad (2.10)$$

Finally, the ρ factors for heavy-heavy currents are defined as

$$\rho_{J_\Gamma}^{hh'} \equiv \frac{Z_{J_\Gamma}^{hh'}}{\sqrt{Z_{V_4}^{hh} Z_{V_4}^{h'h'}}} = 1 + g_0^2 \rho_{J_\Gamma}^{hh'[1]} + \dots \quad (2.11)$$

3. Procedure

In this work we use the automated perturbation theory techniques developed by Lüscher and Weisz [4] to generate the Feynman rules for the lattice actions. We then integrate the loop diagrams by “brute-force” using VEGAS [5]. The advantage of using automated perturbation theory is that it is relatively easy to switch actions [6]. Indeed, we have results for the current renormalizations for two gluon actions, two light quark actions and the heavy quark action.

The one-loop diagrams for the vertex corrections (including the rotations) are given in Ref. [3]. We have performed the following tests of our calculation:

- For the automated perturbation theory code, we have compared our vertices and propagators against known results.
- We have written two independent programs for calculating the current renormalizations based on the automated perturbation theory code.
- We have a third independent calculation of the current renormalizations using traditional semi-analytic methods.
- Our results for the heavy-heavy currents agree with those of Ref. [3] when we switch from the improved gluon propagator to the Wilson gluon propagator. We also reproduce the results of Ref. [3] for heavy-light currents with Clover light quarks and Wilson glue.
- Our result for the Asqtad naive wave function renormalization agrees with Ref. [7], and our result for the naive-naive vertex correction with Wilson glue agrees with Ref. [8].

4. Results




Figure 1: $Z_{V_4}^{hh[1]}$ for equal masses as a function of $m_1^{[0]}$.

Figures 1–4 illustrate our results for the Z 's, ρ 's, and q^* 's as functions of the heavy-quark mass, $m_1^{[0]}a$. The q^* 's are calculated from the log moments using Eqn. (13) of Ref. [9].

Our results for the heavy-heavy currents are very similar to those of Ref. [3], since they differ only in the gluon propagator. The main features of the mass dependence are the same. Figures 1–2 show results for the degenerate mass V_4 current. We also have results for the other currents (V_i , A_4 , A_i) as well as results for currents with unequal masses. In the massless limit we find

$$Z_{V_4}^{hh[1]}(m_1^{[0]} = 0) = -0.10056(3) , \quad (4.1)$$

in good agreement with Ref. [10]. This is another test of our calculation.

Figure 3 shows a comparison of the current renormalization of the heavy-naive A_4 current with the corresponding ρ factor, and Figure 4 shows $\rho_{V_4}^{hl[1]}$ and $\rho_{V_i}^{hl[1]}$. First, the general features of the heavy-quark mass dependence are similar to the results of Ref. [3]. Second, $\rho_{A_4}^{hl[1]}$ is significantly smaller than $Z_{A_4}^{hl[1]}$ over the relevant mass range. Hence, the cancellation between the numerator and denominator of Eq. (2.8) already observed in Ref. [3] also takes place for heavy-naive currents.




Figure 2: $q^* a$ for $Z_{V_4}^{hh[1]}$ for equal masses as a function of $m_1^{[0]}$.

In the massless limit we find:

$$\begin{aligned} \rho_{V_4}^{hl[1]}(m_1^{[0]} = 0) &= -3.038(2) \cdot 10^{-3} \\ \rho_{V_i}^{hl[1]}(m_1^{[0]} = 0) &= -3.05(5) \cdot 10^{-3} \end{aligned} \quad (4.2)$$

We also have results for the naive V_4 current renormalization. In the massless limit we find

$$Z_{V_4}^{ll[1]}(m_0 = 0) = -0.10457(4). \quad (4.3)$$

We have studied the mass dependence of $Z_{V_4}^{ll[1]}$ by varying m_0 between zero and the strange quark mass. We find that $Z_{V_4}^{ll[1]}$ is essentially independent of m_0 .

In summary, we have calculated the current renormalizations relevant for the numerical analyses of heavy-light decay constants and semileptonic form factors performed by the Fermilab Lattice and MILC collaborations. We calculate the full mass dependence of the Z 's and ρ 's. The one-loop corrections to the ρ factors are small. They vary roughly between 0.4% and 4%, depending on lattice spacing.

Acknowledgements

We thank Howard Trotter for his help with testing M.N.'s automated perturbation theory code,




Figure 3: Comparison of $Z_{A4}^{hl[1]}$ with ρ_{A4}^{hl}

in particular by checking our Asqtad vertices and propagators. This work was supported in part by the DOE under grant no. DE-FG02-91ER40677 and by the Junta de Andalucía [P05-FQM-437 and P06-TIC-02302]. The numerical calculations for this work were carried out on the Fermilab lattice QCD clusters, which are a computing resource of the USQCD collaboration and are funded by the DOE. We are grateful to the Fermilab Computing Divisions for operating and maintaining the clusters. Fermilab is operated by Fermi Research Alliance, LLC, under Contract No. DE-AC02-07CH11359 with the United States Department of Energy.

References

- [1] C. Aubin *et al.*, Phys. Rev. **D70**, 114501 (2004); *ibid.* **D70**, 094505 (2004); C. Bernard *et al.*, *ibid.* **D64**, 054506 (2001).
- [2] C. Aubin *et al.*, Phys. Rev. Lett. **95**, 122002 (2005) [hep-lat/0506030]; *ibid.* **94**, 011601 (2005) [hep-ph/0408306]; J. Laiho, these proceedings; P. Mackenzie, these proceedings; J. Simone, these proceedings.
- [3] J. Harada, S. Hashimoto, A. S. Kronfeld and T. Onogi, Phys. Rev. **D65**, 094514 (2002) [hep-lat/0112045]; J. Harada, S. Hashimoto, K. I. Ishikawa, A. S. Kronfeld, T. Onogi and N. Yamada, *ibid.* **D65**, 094513 (2002), [Erratum-*ibid.* **D71**, 019903 (2005)] [hep-lat/0112044].




Figure 4: ρ factors for heavy-naive currents

- [4] M. Lüscher and P. Weisz, Nucl. Phys. **B266**, 309 (1986).
- [5] G. P. Lepage, J. Comp. Phys. **27**, 192 (1978).
- [6] H. D. Trottier, Nucl. Phys. Proc. Suppl. **129**, 142 (2004) [hep-lat/0310044]; M. A. Nobes and H. D. Trottier, Nucl. Phys. Proc. Suppl. **129**, 355 (2004) [hep-lat/0309086].
- [7] E. Gulez, J. Shigemitsu and M. Wingate, Phys. Rev. **D69**, 074501 (2004) [hep-lat/0312017].
- [8] W. Lee and S.R. Sharpe, Phys. Rev. **D66**, 114501 (2002).
- [9] K. Hornbostel, G. P. Lepage and C. Morningstar, Phys. Rev. **D67**, 034023 (2003) [hep-ph/0208224].
- [10] Y. Taniguchi and A. Ukawa, Phys. Rev. **D58**, 114503 (1998) [hep-lat/9806015]; S. Aoki, K. I. Nagai, Y. Taniguchi and A. Ukawa, *ibid.* **D58**, 074505 (1998) [hep-lat/9802034].